

High-Pressure Electrical Resistivity Measurements of EuFe_2As_2 Single Crystals

N Kurita^{1,2}, M Kimata^{1,2}, K Kodama^{1,3}, A Harada¹, M Tomita¹,
H S Suzuki¹, T Matsumoto¹, K Murata⁴, S Uji^{1,2,3} and T Terashima^{1,2}

¹National Institute for Materials Science, Tsukuba, Ibaraki 305-0003, Japan

²JST, Transformative Research-Project on Iron Pnictides (TRIP), Chiyoda, Tokyo 102-0075, Japan

³Graduate School of Pure and Applied Sciences, University of Tsukuba, Ibaraki 305-0003, Japan

⁴Division of Molecular Materials Science, Graduate School of Science, Osaka City University, Osaka 558-8585, Japan

E-mail: KURITA.Nobuyuki@nims.go.jp

Abstract.

High-pressure electrical resistivity measurements up to 3.0 GPa have been performed on EuFe_2As_2 single crystals with residual resistivity ratios $RRR=7$ and 15. At ambient pressure, a magnetic / structural transition related to FeAs-layers is observed at $T_0=190$ K and 194 K for samples with $RRR=7$ and 15, respectively. Application of hydrostatic pressure suppresses T_0 , and then induces similar superconducting behavior in the samples with different RRR values. However, the critical pressure ~ 2.7 GPa, where $T_0 \rightarrow 0$, for the samples with $RRR=15$ is slightly but distinctly larger than ~ 2.5 GPa for the samples with $RRR=7$.

1. Introduction

Since the discovery of superconductivity in $\text{LaFeAs}(\text{O},\text{F})$ with $T_c=26$ K [1], a family of Fe-pnictide superconductors has attracted much attention. In particular, $A\text{Fe}_2\text{As}_2$ ($A=\text{Ca}, \text{Sr}, \text{Ba}, \text{Eu}$, etc.) with a tetragonal ThCr_2Si_2 -type structure has been intensively studied because of the availability of stoichiometric single crystals with high quality. It turned out that, in Fe-pnictide compounds, the superconducting (SC) ground state could appear in accordance with the suppression of a magnetic/structural transition by doping [2]. In the phase diagrams, it is argued that the superconductivity could coexist and/or compete with the antiferromagnetism [3, 4]. However, a random potential introduced by doping could smear the intrinsic SC properties. For understanding the origin of the high- T_c superconductivity with T_c up to 55 K [5], it is of considerable importance to probe the systematic change of ground states using high-quality single crystals. An alternative way to tune the ground state is to apply hydrostatic pressure (P). For instance, recent high- P ac-susceptibility and resistivity measurements have revealed that $A\text{Fe}_2\text{As}_2$ ($A=\text{Sr}, \text{Eu}$) exhibits P -induced bulk superconductivity by suppressing the magnetic/structural transition [6–10]. Meanwhile, superconductivity under hydrostatic P is absent in CaFe_2As_2 [11–14], and remains a controversial issue in BaFe_2As_2 [8, 15–17].

Among the $A\text{Fe}_2\text{As}_2$ series, EuFe_2As_2 is quite unique because the localized Eu^{2+} moments order antiferromagnetically at $T_N \sim 20$ K, in addition to the magnetic/structural transition related to FeAs-layers at $T_0 \sim 190$ K [18–21]. Interestingly, the magnetic order of Eu^{2+} moments

can be detected even in the SC state induced by doping or application of pressure, which could be a main reason for the novel reentrant-SC-like behavior [9, 10, 22–25].

Here, we report the results of high- P electrical resistivity measurements in EuFe_2As_2 using newly grown single crystals with a residual resistivity ratio (RRR) as high as 15. At ambient P , the magnetic/structural transition occurs at a higher temperature of $T_0 = 194$ K, compared with 190 K for single crystals with $RRR = 7$. Consequently, it is found that the higher quality single crystal requires higher- P to suppress T_0 , and to induce the SC ground state in EuFe_2As_2 .

2. Experimental Details

Single crystals of EuFe_2As_2 were grown by Bridgman method from a stoichiometric mixture of the constituent elements. In this study, we examined several crystals from two different batches with residual resistivity ratios $RRR = 7$ and 15, where RRR is defined as $\rho_{300\text{K}}/\rho_{4\text{K}}$. Single crystals measured in Ref [10] were taken from a batch with $RRR = 7$. High-pressure resistivity measurements of samples with $RRR = 7$ and 15 have been performed simultaneously up to 3.0 GPa using a hybrid-type piston cylinder pressure device [26]. The resistivity was measured by the four-probe method with an ac current $I = 0.3$ mA in the ab -plane. To generate hydrostatic pressure, Daphne 7474 (Idemitsu Kosan) oil, which remains in a liquid state up to 3.7 GPa at room temperature [27], was used as a pressure-transmitting medium. Samples were cooled down in Oxford ^4He system, slowly with an average rate of 0.5 K/min. Applied pressure was estimated at 4.2 K from the resistance change of a calibrated Manganin wire [28].

3. Results and Discussions

Figure 1 shows the temperature (T) dependence of electrical resistivity scaled at 300 K ($\rho/\rho_{300\text{K}}$) in EuFe_2As_2 single crystals with $RRR = 7$ and 15, where RRR is determined as $\rho_{300\text{K}}/\rho_{4\text{K}}$. The measurement was performed in zero field at ambient pressure outside a pressure device with current direction $I \parallel ab$. To our knowledge, $RRR = 15$ is the largest value in EuFe_2As_2 single crystals [9, 21, 29]. Overall T -variations of the resistivity in the samples with $RRR = 7$ and 15




Figure 1. (Color online) The scaled electrical resistivity $\rho/\rho_{300\text{K}}$ versus temperature in EuFe_2As_2 single crystals with $RRR = 7$ and 15. The measurement was carried out in zero-field at ambient pressure with the current direction $I \parallel ab$. Upper left and lower right insets represent the expanded views around $T = T_0$ and T_N , respectively. The data for the sample with $RRR = 7$ in the lower right inset is arbitrarily shifted in vertical direction for clarity.

are qualitatively similar to each other, and are consistent with previous results [9, 20, 21, 29]. It is worthwhile to mention that, as shown in the upper left inset, a magnetic/structural transition temperature $T_0 = 194$ K for the sample with $RRR = 15$ is higher than $T_0 = 190$ K for the sample with $RRR = 7$. This would be the reason why samples with $RRR = 15$ needs higher pressure (P) to suppress T_0 , as will be discussed below. The Néel temperature T_N of the localized Eu^{2+} moments for the sample with $RRR = 15$ is slightly higher than the value for the sample with $RRR = 7$, as can be seen in the lower right inset.

Next, we turn to the pressure effect on the electrical resistivity for the samples with $RRR = 7$ (Fig. 2) and 15 (Fig. 3), which are simultaneously measured in the same pressure device. With increasing P , the resistivity peak related to the magnetic/structural transition is suppressed to a lower temperature in both samples as shown in Figs. 2(a) and 3(a). For the sample with $RRR = 7$, a reminiscence of the peak is clearly recognized at 2.38 GPa around 100 K, and faintly visible at 2.46 GPa around 70 K as shown in Fig. 2(a). At 2.55 GPa, there is no detectable anomaly, which implies that the critical pressure P_c , where $T_0 \rightarrow 0$, may be about 2.5 GPa. For the sample with $RRR = 15$, P_c would be ~ 2.7 GPa since the resistivity hump is slightly recognized at 2.69 GPa, but undetectable at 2.77 GPa as shown in Fig. 3(a). It is of interest that the resistivity follows nearly T -linear behavior above T_c at 2.55 and 2.77 GPa ($P \sim P_c$) for samples with $RRR = 7$ and 15, respectively, as guided by a dashed line. A similar T -variation of resistivity was also reported in several optimally-doped Fe-pnictide superconductors [4, 30–32]. For the sample with $RRR = 7$, a resistivity upturn and a small maximum, as indicated by an arrow in Figs. 2(b), in the broad SC transition below 31 K are observed at $P = 2.38$ GPa ($< P_c$). It suggests that the superconductivity is suppressed by the magnetic order of the Eu^{2+} moments; consequently, reentrant-SC-like behavior appears. A similar behavior is also slightly seen for the sample with $RRR = 15$ (Figs. 3(b)), but more smeared out. At $P > P_c$, resistivity exhibits sharp SC transitions to zero-resistivity with $T_c \sim 30$ K for both samples. With increasing P , the SC




Figure 2. ρ vs T of a EuFe_2As_2 single crystal ($RRR = 7$) up to 3.0 GPa in the temperature ranges (a) 30 – 120 K and (b) 15 – 40 K. An arrow in (b) indicates an anomaly attributed to T_N




Figure 3. ρ vs T of a EuFe_2As_2 single crystal ($RRR = 15$) up to 3.0 GPa in the temperature ranges (a) 30 – 120 K and (b) 15 – 40 K. Arrows in (b) indicate anomalies attributed to T_N .

transitions persist up to 3.00 GPa although the T_c continuously decreases. Thus, the P -variation of the resistive behavior between the samples with different quality is qualitatively similar to each other, and is consistent with the previous result [10]. However, $P_c \sim 2.7$ GPa for the sample with $RRR=15$ is slightly but distinctly larger than ~ 2.5 GPa for the sample with $RRR=7$, which may be as a consequence of the larger value of T_0 for the higher-quality sample at ambient- P . We have repeated similar high- P resistivity measurements using several single crystals, and confirmed that the observed difference in the magnitude of T_0 and P_c between the samples with $RRR=7$ and 15 is beyond the error of the pressure estimation ($\pm 2-3 \times 10^{-2}$ GPa) [28]. Another meaningful issue, which probably relates to the sample quality, is the width of a SC transition ΔT_c . The minimum values of ΔT_c are 1 K and 0.8 K for samples with $RRR=7$ and 15, respectively. These facts suggest that the higher-quality single crystals have larger values of T_0 and P_c as well as a sharper SC transition in EuFe_2As_2 .

Until now, there has been no report concerning the quantum oscillation in EuFe_2As_2 , despite the importance for understanding the Fermi surface topology and mass renormalization. In fact, we have already tried de Haas-van Alphen (dHvA) measurements of EuFe_2As_2 using the samples with $RRR=7$ at 0.6 K with fields up to 35 T, but could not detect any dHvA oscillation. Given that quantum oscillations were successfully detected in SrFe_2As_2 ($RRR \sim 8$) [33] and BaFe_2As_2 ($RRR=10$) [34], it is worthwhile to perform the dHvA measurement of EuFe_2As_2 using the newly grown single crystals with $RRR=15$.

4. Conclusions

We have performed high-pressure electrical resistivity measurements up to 3.0 GPa in EuFe_2As_2 single crystals with $RRR=7$ and 15. At ambient pressure, a magnetic/structural transition occurred at $T_0 = 190$ K and 194 K for the samples with $RRR=7$ and 15, respectively. Although P -induced superconductivity was confirmed in the samples with different RRR values, the critical pressure $P_c \sim 2.7$ GPa for the samples with $RRR=15$ was slightly but distinctly larger than ~ 2.5 GPa for the samples with $RRR=7$.

References

- [1] Kamihara Y, Watanabe T, Hirano M and Hosono H 2008 *J. Am. Chem. Soc.* **130** 3296
- [2] Ishida K, Nakai Y and Hosono H 2009 *J. Phys. Soc. Jpn.* **78** 062001 and references therein.
- [3] Chen H, Ren Y, Qiu Y, Bao W, Liu R H, Wu G, Wu T, Xie Y L, Wang X F, Huang Q and Chen X H 2009 *Europhys. Lett.* **85** 17006
- [4] Wang X F, Wu T, Wu G, Liu R H, Chen H, Xie Y L and Chen X H 2009 *New J. Phys.* **11** 045003
- [5] Ren Z A, Lu W, Yang J, Yi W, Shen X L, Li Z C, Che G C, Dong X L, Sun L L, Zhou F and Zhou Z X 2008 *Chin. Phys. Lett.* **25** 2215
- [6] Alireza P L, Ko Y T C, Gillett J, Petrone C M, Cole J M, Lonzarich G G and Sebastian S E 2009 *J. Phys.: Condens. Matter* **21** 012208
- [7] Kotegawa H, Sugawara H and Tou H 2009 *J. Phys. Soc. Jpn.* **78** 013709
- [8] Matsubayashi K, Katayama N, Ohgushi K, Yamada A, Munakata K, Matsumoto T and Uwatoko Y 2009 *J. Phys. Soc. Jpn.* **78** 073706
- [9] Miclea C F, Nicklas M, Jeevan H S, Kasinathan D, Hossain Z, Rosner H, Gegenwart P, Geibel C and Steglich F 2009 *Phys. Rev. B* **79** 212509
- [10] Terashima T, Kimata M, Satsukawa H, Harada A, Hazama K, Uji S, Suzuki H S, Matsumoto T and Murata K 2009 *J. Phys. Soc. Jpn.* **78** 083701
- [11] Torikachvili M S, Bud'ko S L, Ni N and Canfield P C 2008 *Phys. Rev. Lett.* **101** 057006.
- [12] Park T, Park E, Lee H, Klimczuk T, Bauer E D, Ronning F and Thompson J D 2009 *J. Phys.: Condens. Matter* **20** 322204
- [13] Lee H, Park E, Park T, Sidorov V A, Ronning F, Bauer E D and Thompson J D 2009 *Phys. Rev. B* **80** 024519
- [14] Yu W, Aczel A A, Williams T J, Bud'ko S L, Ni N, Canfield P C and Luke G M 2009 *Phys. Rev. B* **79** 020511
- [15] Ishikawa F, Eguchi N, Kodama M, Fujimaki K, Einaga M, Ohmura A, Nakayama A, Mitsuda A and Yamada Y 2009 *Phys. Rev. B* **79** 172506

- [16] Colombier E, Bud'ko S L, Ni N and Canfield P C 2009 *Phys. Rev. B* **80** 224518
- [17] Yamazaki T, Takeshita N, Kobayashi R, Fukazawa H, Kohori Y, Kihou K, Lee C.-H., Kito H, Iyo A and Eisaki H 2010 *Phys. Rev. B* **81** 224511
- [18] Raffius H, Mörsen E, Mosel B D, Müller Warmuth W, Jeitschko W, Terbüchte L and Vomhof T 1993 *J. Phys. Chem. Solids* **54** 135
- [19] Tegel M, Rotter M, Weiss V, Schappacher F M, Pottgen R and Johrendt D 2008 *J. Phys.: Condens. Matter* **20** 452201
- [20] Ren Z, Zhu Z W, Jiang S, Xu X F, Tao Q, Wang C, Feng C M, Cao G H and Xu Z A 2008 *Phys. Rev. B* **78** 052501
- [21] Jeevan H S, Hossain Z, Kasinathan D, Rosner H, Geibel C and Gegenwart P 2008 *Phys. Rev. B* **78** 052502
- [22] Ren Z, Lin X, Tao Q, Jiang S, Zhu Z, Wang C, Cao G and Xu Z 2008 *Phys. Rev. B* **78** 092406.
- [23] Jiang S, Xing H, Xuan G, Ren Z, Wang C, Xu Z and Cao G 2009 *Phys. Rev. B* **80** 184514.
- [24] Ren Z, Tao Q, Jiang S, Feng C, Wang C, Dai J, Cao G and Xu Z 2009 *Phys. Rev. Lett.* **102** 137002.
- [25] Zheng Q J, He Y, Wu T, Wu G, Chen H, Ying J J, Liu R H, Wang X F, Xie Y L, Yan Y J, Li Q J and Chen X H 2009 arXiv:0907.5547v1
- [26] Uwatoko Y, Hedo M, Kurita N, Koeda M, Abliz M and Matsumoto T 2003 *Physica C* **329-333** 1658
- [27] Murata K, Yokogawa K, Yoshino H, Klotz S, Munsch P, Irizawa A, Nishiyama M, Iizuka K, Nanba T, Okada T, Shiraga Y and Aoyama S 2008 *Rev. Sci. Instrum.* **79** 085101
- [28] Terashima T, Tomita M, Kimata M, Satsukawa H, Harada A, Hazama K, Uji S, Suzuki H S, Matsumoto T and Murata K 2009 *J. Phys. Soc. Jpn.* **78** 118001
- [29] Mitsuda A, Matoba T, Wada H, Ishikawa F and Yamada Y 2010 *J. Phys. Soc. Jpn.* **79** 073704
- [30] Liu R H, Wu G, Wu T, Fang D F, Chen H, Li S Y, Liu K, Xie Y L, Wang X F, Yang R L, Ding L, He C, Feng D L and Chen X H 2008 *Phys. Rev. Lett.* **101** 087001
- [31] Gooch M, Bing L, Lorenz B, Guloy A M and Chu C W 2009 *Phys. Rev. B* **79** 104504
- [32] Jiang S, Xing H, Xuan G, Wang C, Ren Z, Feng C, Dai J, Xu Z and Cao G 2009 *J. Phys.: Condens. Matter* **21** 382203
- [33] Sebastian S E, Gillett J, Harrison N, Lau P H C, Singh D J, Mielke C H and Lonzarich G G 2008 *J. Phys.: Condens. Matter* **20** 422203
- [34] Analytis J G, McDonald R D, Chu J -H, Riggs S C, Bangura A F, Kucharczyk C, Johannes M and Fisher I R 2009 *Phys. Rev. B* **80** 064507